arXiv:1207.6946v1 [nlin.SI] 30 Jul 2012

Application of the modified method of simplest equation for obtaining exact traveling-wave solutions for the extended Korteweg - de Vries equation and generalized Camassa-Holm equation

Nikolay K. Vitanov[1,2]*, Zlatinka I. Dimitrova[3], Holger Kantz [2]

[1] Institute of Mechanics, Bulgarian Academy of Sciences, Akad. G. Bonchev Str., Bl. 4, 1113 Sofia, Bulgaria
[2] Max-Planck-Institute for the Physics of Complex Systems, Noethnitzerstr. 38, 01187 Dresden, Germany
[3] "G. Nadjakov" Institute for Solid State Physics, Bulgarian Academy of Science, Blvd. Tzarigradsko Chaussee 72, 1784 Sofia, Bulgaria

Abstract

The modified method of simplest equation is applied to the extended Korteweg - de Vries equation and to generalized Camassa - Holm equation. Exact traveling wave solutions of these two nonlinear partial differential equations are obtained. The equations of Bernoulli, Riccati and the extended tanh - equation are used as simplest equations. Some of the obtained solutions correspond to surface water waves.

MSC 2010 numbers:

35Q35 : PDEs in connection with fluid mechanics
35Q53 : KdV-like equations

Key words:

method of simplest equation, exact traveling-wave solutions, generalized Camassa - Holm equation, extended Korteweg - de Vries equation, water waves

---

*e-mail: vitanov@imbm.bas.bg; vitanov@mpipks-dresden.mpg.de

1 Introduction

Because of the nonlinearity of the Navier - Stokes equation, Boussinesq equations, etc., the hydrodynamics is a rich sorce of nonlinear problems [1] - [9]. In addition in the last decades the nonlinear partial differential equations are widely used for modeling natural and social phenomena and systems. Examples are [10, 11] from neurology, [12] -[15] from the theory of solitons, [16] from the biology, [17, 18] from theory of dynamical systems, chaos theory and ecology, and [19] - [26] from hydrodynamics and theory of turbulence. The knowledge of exact solutions of the model PDEs is usefull in several directions. First of all such solutions describe different kinds of waves. In addition the exact solutions are useful as initial condition for computer programs that simulate processes in the investigated system. Because of this an important research area is connected to obtaining exact analytical or approximate numerical solutions of such model PDEs. Among the most famous methods for obtaining solutions of exact solvable equations are the inverse scattering transform [27] - [30] and the method of Hirota [31]. In the course of years numerous approaches for obtaining exact special solutions of nonlinear PDE have been developed [32] - [39]. By means of such methods numerous exact solutions of many equations have been obtained such as for an example the Kuramoto-Shivasinsky equation [37], equations, connected to the models of population dynamics [40] - [52], sine - Gordon equation [53] - [60], sinh-Gordon or Poisson - Boltzmann equation [61] - [63], Lorenz - like systems [64], etc.

One of the dirct methods for obtaining exact and approximate solutions of nonlinear partial differential equations is the method of simplest equation [37] - [39]. This method ant its modification called modified method of simplest equation have been applied to numerous nonlinear PDEs [65], [66], such as Fisher equation and Fisher - like equations [67], classes of reaction - diffusion and reaction - telegraph equations [50], generalized Kuramoto - Sivashinsky equation [38], generalized Swift - Hohenberg and generalized Rayleigh equations [39], generalized Deagsperis - Processi equation and b-equation [68]. In this paper we shall apply the modified method of simplest equation to the extended Korteweg - de Vries equation and to generalized Camassa - Holm equation. The goal of the paper is to obtain exact traveling - wave solutions to the last two equations.

The paper is organized as follows. In Sect. 2 information from the theory of water waves is presented which connects the Navier - Stokes equation and the profile of a class of water waves to the exact traveling - wave solutions of the extended Korteweg - de Vries equation and Camassa - Holm equation. Sect. 3 contains information about the modified method of simplest equation needed for obtaining exact traveling - wave solution of the extended Korteweg - de Vries equation and of the generalized Camassa - Holm equation. In sect 4. we obtain exact traveling-wave solutions of the extended Korteweg - de Vries equation on the basis of the use of the equations of Bernoulli, Riccati and extended tanh - equation as simplest equations. In section 5 using the same procedure we obtain exact traveling - wave solutions of the generalized Camassa - Holm equation. Several concluding remarks are summarized in Sect. 6.

2 Water waves, extended Kortweg - de - Vries equation and generalized Camassa - Holm equation

The research on water waves is a rich sorce of nonlinear problems [69] - [71]. We shall obtain the extended Korteweg - de Vries equation and the Camassa - Holm equation following Johnson [72]. Let us consider incompressible inviscid fluid with zero surface tension. The fluid moves over an steady intermediate bed with coordinate $z = z^*(x, y)$. The pressure on the top boundary of the fluid is constant: $P = P_a$, where P_a is the atmospheric pressure. The system of model equations for description of the movement of the fluid consists of the Navier - Stokes equation and continuity equation

$$\frac{\partial \vec{u}}{\partial t} + \vec{u} \cdot \nabla \vec{u} = -\frac{1}{\rho} \nabla P + \vec{F} \tag{2.1}$$

$$\nabla \cdot \vec{u} = 0 \tag{2.2}$$

where $\vec{u} = (u, v, w)$ is the velocity of the fluid, ρ is the density of the fluid and the acting force is gravity: $\vec{F} = (0, 0, -g)$. The fluid surface is at $z = h(x, y, t)$ and the boundary conditions are as follows.

- On the fluid surface

$$P = P_a; \quad w = \frac{\partial h}{\partial t} + (\vec{u} \cdot \nabla) h \tag{2.3}$$

- On the bed $z = z^*(x, y)$

$$w = u \frac{\partial z^*}{\partial x} + v \frac{\partial z^*}{\partial y} \tag{2.4}$$

Let h_0 be the undisturbed depth of the water. Then we introduce the non-dimensional deviation η from h_0 as follows

$$h = h_0 + a\eta \tag{2.5}$$

In addition we introduce the non - dimensional pressure p

$$P = P_a + \rho g (h_0 - z) + \rho g h_0 p \tag{2.6}$$

p measures the deviation from the hydrostatic pressure. Our goal is to study wave propagation on the fluid surface. The scale of this motion is the typical wavelegth of the surface wave λ. We shall use λ as unit for length.

Next we perform a non-dimensionalization of the model equations by the following substitutions (→ means "replace by" and in this way the notation is retained)

$$x \to \lambda x; \quad y \to \lambda y; \quad z \to h_0 z; \quad t \to \frac{\lambda}{\sqrt{g h_0} t};$$
$$u \to \sqrt{g h_0} u; \quad v \to \sqrt{g h_0} v; \quad w \to \frac{h_0 \sqrt{g h_0}}{\lambda} w; \quad z^* \to h_0 z^* \tag{2.7}$$

In addition we introduce two important parameters: amplitude parameter ϵ and shallowness parameter δ:

$$\epsilon = \frac{a}{h_0}; \quad \delta = \frac{h_0}{\lambda} \tag{2.8}$$

Further we rescale the non-dimensionalized equations with respect to the amplitude parameter ϵ:

$$u \to \epsilon u; \quad v \to \epsilon v; \quad w \to \epsilon w; \quad p \to \epsilon p \tag{2.9}$$

and obtain the system of equations

$$\frac{\partial u}{\partial t} + \epsilon(\vec{u} \cdot \nabla)u = -\frac{\partial p}{\partial x} \tag{2.10}$$

$$\frac{\partial v}{\partial t} + \epsilon(\vec{u} \cdot \nabla)v = -\frac{\partial p}{\partial y} \tag{2.11}$$

$$\nabla \cdot \mathbf{u} = 0 \tag{2.12}$$

with boundary conditions:

$$p = \eta; \quad w = \frac{\partial \eta}{\partial t} + \epsilon \left(u\frac{\partial \eta}{\partial x} + v\frac{\partial \eta}{\partial y} \right) \tag{2.13}$$

on $z = 1 + \epsilon\eta$ and

$$w = u\frac{\partial z^*}{\partial x} + v\frac{\partial z^*}{\partial y} \tag{2.14}$$

on $z = z^*$. We continue the scaling of (2.10) - (2.12) as follows: $\zeta = \sqrt{\epsilon}(x - t)$; $\tau = \epsilon\sqrt{\epsilon}t$; $w = \sqrt{\epsilon}W$ and expand all quantities

$$q \sim \sum_{n=0}^{\infty}\sum_{m=0}^{\infty} \epsilon^n \delta^{2m} q_{nm} \tag{2.15}$$

Assuming that $\epsilon \to 0$ and $\delta \to 0$ (small amplitude water waves) from (2.10) - (2.12) we obtain that the non-dimensionalized height of the water surface $\eta = \eta_{00} + \epsilon_1\eta_{10} + \delta^2\eta_{01} + \epsilon\delta^2\eta_{11}$ satisifies the higher order Korteweg - de Vries equation

$$2\frac{\partial \eta}{\partial \tau} + 3\eta\frac{\partial \eta}{\partial \zeta} + \frac{1}{3}\delta^2\frac{\partial^3 \eta}{\partial \zeta^3} - \frac{3}{4}\epsilon\eta^2\frac{\partial \eta}{\partial \zeta} = -\frac{1}{12}\epsilon\delta^2\left(23\frac{\partial \eta}{\partial \zeta}\frac{\partial^2 \eta}{\partial \zeta^2} + 10\eta\frac{\partial^3 \eta}{\partial \zeta^3}\right) + O(\epsilon^2, \delta^4) \tag{2.16}$$

and to the same order the solution is

$$u \approx \eta - \frac{1}{4}\eta^2 + \epsilon\delta^2\left(\frac{1}{3} - \frac{1}{2}z^2\right)\frac{\partial^2 \eta}{\partial \zeta^2} \tag{2.17}$$

where z is connected to the depth of the moving fluid. Eq. (2.16) is the extended Korteweg - de Vries equation and it will be one of equations we shall discuss below.

Another important water wave equation - the Camassa - Holm equation can be obtained in similar way like (2.16) above. The Camassa - Holm equation describes the $x-$ component u of the fluid velocity at certain depth z_0 below the fluid surface. Cammassa - Holm equation was introduced as completely integrable bi-Hamiltonian dispersive shallow water equation [73]. In the framework discussed here and in [72] the Camassa - Holm equation is

$$\frac{\partial U}{\partial T} + 2\kappa\frac{\partial U}{\partial X} + 3\epsilon U\frac{\partial U}{\partial X} - \epsilon\delta^2\frac{\partial^3 U}{\partial X^2 \partial T} = \epsilon^2\delta^2\left(2\frac{\partial U}{\partial X}\frac{\partial^2 U}{\partial X^2} + U\frac{\partial^3 U}{\partial X^3}\right) \tag{2.18}$$

The error of this equation is $O(\epsilon^3, \epsilon\delta^4)$; it is valid for $z_0 = \frac{1}{\sqrt{2}}$. In addition $\kappa = \frac{4}{5}\sqrt{\frac{3}{5}}$ and the solution of the Camassa - Holm equation equation corresponds within the approximation $O(\epsilon^3, \epsilon\delta^4)$ to the surface wave

$$\eta \sim \sqrt{\frac{5}{3}}\left(U + \frac{1}{4}\sqrt{\frac{5}{3}}\epsilon U^2 - \frac{1}{5}\epsilon\delta^2 \frac{\partial^2 U}{\partial X^2}\right) \tag{2.19}$$

where $T = t\sqrt{\epsilon}$; $X = 2\sqrt{\frac{5}{3}}\left(Z - \frac{3}{5}T\right)$; $Z = x\sqrt{\epsilon}$.

In this paper we shall discuss the following generalized Camassa - Holm equation

$$\begin{aligned} &\frac{\partial U}{\partial T} + p_1 \frac{\partial U}{\partial X} - p_2 \frac{\partial^3 U}{\partial X^2 \partial T} - p_3 \frac{\partial^3}{\partial X^3} f(U) + \\ &p_4 \frac{\partial}{\partial X}\left[\frac{1}{2} g(U) + \frac{p_5}{2}\frac{d^2 f}{dU^2}\left(\frac{\partial U}{\partial X}\right)^2\right] = 0 \end{aligned} \tag{2.20}$$

where $p_1; p_2; p_3; p_4; p_5$ are parameters. Let $f(U) = \frac{1}{2}U^2$; $g(U) = C + 3U^2$ (C is a constant) and $p_3 = p_4 p_5$. Then from Eq. (2.20) we obtain

$$\frac{\partial U}{\partial T} + p_1 \frac{\partial U}{\partial X} - p_2 \frac{\partial^3 U}{\partial X^2 \partial T} + 3p_4 U \frac{\partial U}{\partial X} - 2p_3 \frac{\partial U}{\partial X}\frac{\partial^2 U}{\partial X^2} - p_3 U \frac{\partial^3 U}{\partial X^3} = 0 \tag{2.21}$$

Eq. (2.20) contains the Camassa - Holm equation as particular case when $p_2 = 2\kappa$; $p_4 = \epsilon$; $p_2 = \epsilon\delta^2$; $p_3 = \epsilon^2\delta^2$. In general in Eq. (2.20) $f(U)$ and $g(U)$ can be arbitrary polynomials of U. Finally we note that if we do not set $g(U)$ and all other settings are as for obtaining Eq. (2.21) from Eq. (2.20) we obtain the following generalization of the Camassa - Holm equation

$$\frac{\partial U}{\partial T} + p_1 \frac{\partial U}{\partial X} + \frac{p_4}{2}\frac{\partial}{\partial X} g(U) - p_2 \frac{\partial^3 U}{\partial X^2 \partial T} - 2p_3 \frac{\partial U}{\partial X}\frac{\partial^2 U}{\partial X^2} - p_3 U \frac{\partial^3 U}{\partial X^3} = 0 \tag{2.22}$$

Let $p_2 = p_4 = 1$;$p_1 = p_3$. Then we obtain the version of the generalized Camassa - Holm equation discussed by Tian and Wang [74] (see also [75, 76]).

3 The modified method of simplest equation

The method of simplest equation has been developed by Kudryashov [37, 49] on the basis of a procedure analogous to the first step of the test for the Painleve property. In the modified method of the simplest equation [38, 39] this procedure is substituted by the concept for the balance equation (for details about the balance equation see the subsection of this section). The essence of the method is as follows. Let us have a partial differential equation and let by means of an appropriate ansatz this equation is reduced to a nonlinear ordinary differential equation

$$P\left(u(\xi), \frac{du}{d\xi}, \frac{d^2u}{d\xi^2}, \dots\right) = 0 \tag{3.1}$$

For large class of equations from the kind (3.1) exact solution can be constructed as finite series

$$u(\xi) = \sum_{\mu=-\nu}^{\nu_1} \theta_\mu [V(\xi)]^\mu \tag{3.2}$$

where $\nu > 0$, $\mu > 0$, θ_μ is a parameter and $V(\xi)$ is a solution of some ordinary differential equation referred to as the simplest equation. The simplest equation is of lesser order than (3.1) and we know the general solution of the simplest equation or we know at least exact analytical particular solution(s) of the simplest equation [65, 66].

The modified method of simplest equations can be applied to nonlinear partial differential equations of the kind

$$F\left(\frac{\partial^{\omega_1} u}{\partial x^{\omega_1}}, \frac{\partial^{\omega_2} u}{\partial t^{\omega_2}}, \frac{\partial^{\omega_3} u}{\partial x^{\omega_4} \partial t^{\omega_5}}\right) = G(u) \tag{3.3}$$

where $\omega_3 = \omega_4 + \omega_5$ and

1. $\frac{\partial^{\omega_1} u}{\partial x^{\omega_1}}$ denotes the set of derivatives
$$\frac{\partial^{\omega_1} u}{\partial x^{\omega_1}} = \left(\frac{\partial u}{\partial x}, \frac{\partial^2 u}{\partial x^2}, \frac{\partial u^3}{\partial x^3}, \dots\right)$$
2. $\frac{\partial^{\omega_2} u}{\partial t^{\omega_2}}$ denotes the set of derivatives
$$\frac{\partial^{\omega_2} u}{\partial t^{\omega_2}} = \left(\frac{\partial u}{\partial t}, \frac{\partial^2 u}{\partial t^2}, \frac{\partial u^3}{\partial t^3}, \dots\right)$$
3. $\frac{\partial^{\omega_3} u}{\partial x^{\omega_4} \partial t^{\omega_5}}$ denotes the set of derivatives
$$\frac{\partial^{\omega_3} u}{\partial x^{\omega_4} \partial t^{\omega_5}} = \left(\frac{\partial^2 u}{\partial x \partial t}, \frac{\partial^3 u}{\partial x^2 \partial t}, \frac{\partial u^3}{\partial x \partial t^2}, \dots\right)$$
4. $G(u)$ can be
   - (a) polynomial of u or
   - (b) function of u which can be reduced to polynomial of u by means of Taylor series for small values of u
5. F can be an arbitrary sum of products of arbitrary number of its arguments. Each argument in each product can have arbitrary power. Each of the products can be multiplied by a function of u which can be
   - (a) polynomial of u or
   - (b) function of u which can be reduced to polynomial of u by means of Taylor series for small values of u

For this class of equations the modified method of simplest equation allows us in principle to search for

1. Exact traveling-wave solutions of (3.3) if $G(u)$ and the multiplication functions form item 5. above are polynomials
2. Approximate traveling-wave solutions for small u in all other cases

We are interested in traveling wave solutions

$$u(x,t) = u(\xi) = u(x - vt) \tag{3.4}$$

of the extended Korteweg - de vries equation and of the generalized Camassa - Holm equation. These solutions will be obtained on the basis of the modified method of simplest equation.

In what follows below we shall set $\nu = 0$ and we shall use as simplest equation particular cases of the following equation

$$\frac{dV}{d\xi} = \sum_{\alpha=0}^{\beta} \gamma_\alpha [V(\xi)]^\alpha \tag{3.5}$$

where γ_α is a parameter. Two particular cases of (3.5) are the equations of Bernoulli and Riccati which are well known nonlinear ordinary differential equations and their solutions can be expressed by elementary functions. For the Bernoulli equation $\gamma_0 = 0$; $\gamma_1 = a$; $\gamma_2 = \dots = \gamma_{k-1} = 0$; $\gamma_k = c$; $\gamma_{k+1} = \dots = \gamma_\beta = 0$. The result is

$$\frac{dV}{d\xi} = aV(\xi) + c[V(\xi)]^k \tag{3.6}$$

where k is an integer and $k > 1$. For (3.6) we shall use the solutions

$$V(\xi) = \left\{ \frac{a \exp[a(k-1)(\xi + \xi_0)]}{1 - c\exp[a(k-1)(\xi + \xi_0)]} \right\}^{\frac{1}{k-1}} \tag{3.7}$$

for the case $c < 0$, $a > 0$ and

$$V(\xi) = \left\{ -\frac{a \exp[a(k-1)(\xi + \xi_0)]}{1 + c\exp[a(k-1)(\xi + \xi_0)]} \right\}^{\frac{1}{k-1}} \tag{3.8}$$

for the case $c > 0$, $a < 0$. Above ξ_0 is a constant of integration. For the Riccati equation $\gamma_0 = d$; $\gamma_1 = c$; $\gamma_2 = a$, $\gamma_3 = \dots = \gamma_\beta = 0$. The result is

$$\frac{dV}{d\xi} = a[V(\xi)]^2 + cV(\xi) + d \tag{3.9}$$

We shall use two solutions of (3.9). By a substitution in (3.9) it can be shown that

$$V_1(\xi) = -\frac{c}{2a} - \frac{\theta}{2a} \tanh\left[\frac{\theta(\xi + \xi_0)}{2}\right] \tag{3.10}$$

is a particular solution of the Riccati equation. On the basis of (3.10) we can construct more complicated solution of the Riccati equation which has the form

$$V_2(\xi) = V_1(\xi) + \frac{1}{V_3(\xi)}$$

where $V_3(\xi)$ is solution of the linear equation

$$\frac{dV_3}{d\xi} + [c + 2aV_1(\xi)]V_3(\xi) = -a$$

For the solution $V_2(\xi)$ we obtain

$$V_2(\xi) = -\frac{c}{2a} - \frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right) + \frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta} + 2C^*\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]} \tag{3.11}$$

In (3.10) and (3.11) $\theta^2 = c^2 - 4ad > 0$. In addition ξ_0 and C^* are constants of integration. We shall use the solutions $V_1(\xi)$ and $V_2(\xi)$ below.

As particular case of the use of equation of Riccati as simplest equation we shall consider also the so called extended tanh-function equation

$$\frac{dV}{d\xi} = \overline{d}^2 - \overline{a}^2 V^2 \tag{3.12}$$

used by Fan [34] as basis for the extended tanh-function method. (3.12) is obtained from (3.9) when $b = 0$, $a = -\overline{a}^2$, and $d = \overline{d}^2$. The solution of (3.12) we shall use below is

$$V(\xi) = \frac{\overline{d}}{\overline{a}}\tanh[\overline{a}\,\overline{d}(\xi+\xi_0)] \tag{3.13}$$

where $\overline{a}^2 V(\xi)^2 < \overline{d}^2$ and ξ_0 is a constant of integration. The maximum power of $V(\xi)$ in (3.12) is the same as the maximum power of $V(\xi)$ in (3.9). Because of this the obtained below balance equations for the equation of Riccati and for the extended tanh-equation will be the same. The difference between the two simplest equations is that the equation of Riccati will lead to more complicated exact solutions of the corresponding PDE.

3.1 Balance equations

Let us search for a solution of (3.1) which is of the kind (3.2) with $\eta > \nu$ and where $V(\xi)$ is a solution of (3.5). The substitution of (3.2) and the corresponding simplest equation in (3.1) leads to a polynomial equation of the kind

$$P = \kappa_r V^r + \kappa_{r-1} V^{r-1} + \dots + \kappa_0 = 0 \tag{3.14}$$

where r is some integer and the coefficients κ depend on the parameters of the solved equation as well as on the parameters of the solution. The solution of (3.14) is obtained when all coefficients in (3.14) are equal to 0. This leads to a system of nonlinear algebraic relationships

$$\kappa_l = 0, \quad l = r, r-1, \dots, 0 \tag{3.15}$$

In the classic version of the simplest equation method Kudryashov determines ν_1 by a procedure analogous to the first step of the test for the Painleve property [49]. Below we shall follow another idea [38, 39]. We have to ensure that there are at least two terms containing V^r in (3.14). In order to achieve this we have to balance at least two of the largest powers of the polynomials arising from the terms of (3.1). Below we shall do this for the extended Korteweg - de Vries equation and for the generakized Camassa - Holm equation.

Let us now conseder the extended Kortweg - de vries equation (2.16) As first we shall use the equation of Bernoulli (3.6) as simplest equation. The solution of Eq. (2.16) will be constructed by means of solutions of Eq. (3.6) on the basis of (3.2) with $\nu = 0$. We shall solve generalized version of the extended Korteweg - de Vries equation

$$p_1 \frac{\partial \eta}{\partial \tau} + p_2 \eta \frac{\partial \eta}{\partial \zeta} + p_3 \frac{\partial^3 \eta}{\partial \zeta^3} + p_4 \eta^2 \frac{\partial \eta}{\partial \zeta} + p_5 \frac{\partial \eta}{\partial \zeta} \frac{\partial^2 \eta}{\partial \zeta^2} + p_6 \eta \frac{\partial^3 \eta}{\partial \zeta^3} = 0 \tag{3.16}$$

Eq (3.16) is reduced to Eq. (2.16) for the following values of the parameters

$$p_1 = 2; \;\; p_2 = 3; \;\; p_3 = \frac{1}{3}\delta^2; \;\; p_4 = -\frac{3}{4}\epsilon; \;\; p_5 = \frac{23}{12}\epsilon\delta^2; \;\; p_6 = \frac{5}{6}\epsilon\delta^2 \tag{3.17}$$

We shall introduce the coordinate $\xi = \zeta - v\tau$ and shall search for traveling-wave solutions of Eq. (3.16). In the original coordinates x and t : $\xi = \sqrt{\epsilon}(x - \epsilon v t)$. The application of the modified method of simplest equation to Eq. (3.16) on the basis of equation of Bernoulli as simplest equation leads to the following balance equation

$$\nu_1 = 2(k-1) \tag{3.18}$$

Let now the equation of Riccati (3.9) be used as simplest equation. We again search solution of (3.16) in the form (3.2) with $\nu = 0$. For this case the balance equation is

$$\nu_1 = 2 \tag{3.19}$$

The use of the extended tanh - equation as simplest equation leads to the same balance equation as for the case of Riccati equation namely (3.19).

Now let us discuss the generalized Camassa - Holm equation (2.20). We are interested in traveling wave solutions. Because of this we introduce the coordinate $\xi = X - vT$ where v is the velocity of the corresponding wave. First let the equation of Bernoulli (3.6) be used as simplest equation and the solution of Eq. (2.20) be searched in the form of Eq. (3.2) with $\nu = 0$. The determination of the maximum powers of the terms in Eq. (2.20) leads to the following balance equation

$$\nu_1 = \frac{2(k-1)}{J-I} \tag{3.20}$$

where $k > 1$ and $J > I > 1$. For $I > 1$ another balance equation is possible namely

$$I\nu_1 + 3(k-1) = I\nu_1 + 3(k-1) \tag{3.21}$$

when $(J-I)\nu_1 < 2(k-1)$. This balance is possible as there are two terms in (2.20) which lead to maximum power $I\nu_1 + 3(k-1)$ of terms of the resulting polynomial and these powers are maximum powers for the entire polynomial when $(J-I)\nu_1 < 2(k-1)$.

Finally if $I = 1$ the balance equation is

$$\nu_1 = \frac{2(k-1)}{J-1} \tag{3.22}$$

Another possibility is to use the equation of Riccati (3.9) as simplest equation. The solution of Eq. (2.20) is searched again the form of Eq. (3.2). Several balance equations are possible for this case. They are

$$\nu_1 = I; \quad J < \nu_1 + 2; \quad J < I + 2 \tag{3.23}$$

$$\nu_1 + 2 = J; \quad I < \nu_1; \quad I + 2 < J \tag{3.24}$$

$$J = I + 2; \quad \nu_1 < I; \quad \nu_1 + 2 < J \tag{3.25}$$

The possible balance equations for the case when the extended tanh - equation is used as simplest equation are the same as for the case of Riccati equation.

Finally we would like to note two of the papers of Kudryashov [77, 78] who advises us to be careful when use the methods for obtaining of exact solutions of the nonlinear differential equations. Especially we have to check if the newly obtained solutions can be reduced to some of the already known exact solutions of the correspondent PDE.

4 Exact traveling wave solutions of the extended Korteweg - de Vries equation (2.16)

First of all we shall use the equation of Bernoulli as simplest equation. Let $k = 2$. Then from Eq. (3.18) $\nu_1 = 2$. The subsitution of Eq. (3.2) and Eq. (3.6) in Eq. (3.16) leads to the following system of 7 nonlinear algebraic relationships between the parameters of Eq.

(3.16) and parameters of its solution

$$\begin{aligned}
0 &= c\theta_2^2[6p_5c^2 + p_4\theta_2 + 12p_6c^2] \\
0 &= \theta_2[24p_6\theta_1c^3 + p_6(6\theta_1c^3 + 54\theta_2ac^2) + 6p_5(\theta_1c + 2\theta_2a)c^2 + \\
&\quad 2p_5c(10\theta_2ac + 2\theta_1c^2) + 4p_4\theta_1\theta_2c + p_4\theta_2(\theta_1c + 2\theta_2a)] \\
0 &= 24p_3\theta_2c^3 + 6p_5\theta_1a\theta_2c^2 + p_5(\theta_1c + 2\theta_2a)(10\theta_2ac + 2\theta_1c^2) + \\
&\quad 2p_5\theta_2c(3\theta_1ac + 4\theta_2a^2) + 2p_2\theta_2^2c + 2p_4(2\theta_0\theta_2 + \theta_1^2)\theta_2c + \\
&\quad 2p_4\theta_1\theta_2(\theta_1c + 2\theta_2a) + p_4\theta_2^2\theta_1a + 24p_6\theta_0\theta_2c^3 + p_6\theta_1(6\theta_1c^3 + 54\theta_2ac^2) + \\
&\quad p_6\theta_2(12\theta_1c^2a + 38\theta_2a^2c) \\
0 &= 4p_4\theta_0\theta_1\theta_2c + p_4(2\theta_0\theta_2 + \theta_1^2)(\theta_1c + 2\theta_2a) + 2p_4\theta_1^2\theta_2a + \\
&\quad 2p_2\theta_1\theta_2c + p_2\theta_2(\theta_1c + 2\theta_2a) + p_3(6\theta_1c^3 + 54\theta_2ac^2) + \\
&\quad p_6\theta_0(6\theta_1c^3 + 54\theta_2ac^2) + p_6\theta_1(12\theta_1c^2a + 38\theta_2a^2c) + \\
&\quad p_6\theta_2(7\theta_1a^2c + 8\theta_2a^3) + p_5\theta_1a(10\theta_2ac + 2\theta_1c^2) + \\
&\quad p_5(\theta_1c + 2\theta_2a)(3\theta_1ac + 4\theta_2a^2) + 2p_5\theta_2c\theta_1a^2 \\
0 &= -2vp_1\theta_2c + p_6\theta_0(12\theta_1c^2a + 38\theta_2a^2c) + p_6\theta_1(7\theta_1a^2c + 8\theta_2a^3) + \\
&\quad p_6\theta_2\theta_1a^3 + 2p_2\theta_0\theta_2c + p_2\theta_1(\theta_1c + 2\theta_2a) + p_2\theta_2\theta_1a + \\
&\quad p_3(12\theta_1c^2a + 38\theta_2a^2c) + 2p_4\theta_0^2\theta_2c + 2p_4\theta_0\theta_1(\theta_1c + 2\theta_2a) + \\
&\quad p_4(2\theta_0\theta_2\theta_1^2)\theta_1a + p_5\theta_1a(3\theta_1ac + 4\theta_2a^2) + p_5(\theta_1c + 2\theta_2a)\theta_1a^2 \\
0 &= -vp_1(\theta_1c + 2\theta_2a) + p_4\theta_0^2(\theta_1c + 2\theta_2a) + 2p_4\theta_0\theta_1^2a + p_5\theta_1^2a^3 + \\
&\quad p_2\theta_0(\theta_1c + 2\theta_2a) + p_2\theta_1^2a + p_3(7\theta_1a^2c + 8\theta_2a^3) + p_6\theta_0(7\theta_1a^2c + 8\theta_2a^3) + \\
&\quad p_6\theta_1^2a^3 \\
0 &= \theta_1[-vp_1a + p_4\theta_0^2a + p_6\theta_0a^3 + p_2\theta_0a + p_3a^3]
\end{aligned} \tag{4.1}$$

One solution of this system of algebraic equations is

$$\begin{aligned}
\theta_2 &= -\frac{6c^2(p_5 + 2p_6)}{p_4}; \quad \theta_1 = -\frac{6ac(p_5 + 2p_6)}{p_4} \\
\theta_0 &= -\frac{-2p_3p_4 + a^2p_5^2 + 3a^2p_5p_6 + p_2p_5 + 2p_2p_6 + 2a^2p_6^2}{2p_4(p_5 + p_6)} \\
v &= \frac{a^4p_5^4 + 4p_3^2p_4^2 - 4p_2p_3p_4p_6 - p_2^2p_5^2 - 2p_2^2p_5p_6 + 4a^4p_6p_5^3 + 5a^4p_6^2p_5^2 + 2a^4p_6^3p_5}{4p_1p_4(p_5 + p_6)^2}
\end{aligned} \tag{4.2}$$

The solution of Eq. (3.16) is

$$\begin{aligned}
\eta = &-\frac{-2p_3p_4 + a^2p_5^2 + 3a^2p_5p_6 + p_2p_5 + 2p_2p_6 + 2a^2p_6^2}{2p_4(p_5 + p_6)} - \\
&\frac{6a^2c(p_5 + 2p_6)}{p_4}\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right]\left\{1 + c\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right]\right\}
\end{aligned} \tag{4.3}$$

for the case $a > 0$, $c < 0$ and

$$\eta = -\frac{-2p_3p_4 + a^2p_5^2 + 3a^2p_5p_6 + p_2p_5 + 2p_2p_6 + 2a^2p_6^2}{2p_4(p_5 + p_6)} + \frac{6a^2c(p_5 + 2p_6)}{p_4} \left[\frac{\exp[a(\xi + \xi_0)]}{1 + c\exp[a(\xi + \xi_0)]}\right] \left\{1 - c\left[\frac{\exp[a(\xi + \xi_0)]}{1 + c\exp[a(\xi + \xi_0)]}\right]\right\} \tag{4.4}$$

for the case $a < 0$, $c > 0$. Let us discuss the case $a > 0$; $c < 0$. From Eq. (3.17) we obtain the following solution of the extended Korteweg - de Vries equation (2.16)

$$\eta = \frac{540 + 473\epsilon\delta^2a^2}{198\epsilon} + \frac{86\delta^2ca^2}{3} \left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right] \left\{1 + c\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right]\right\} \tag{4.5}$$

For the surface wave from Eq. (2.17) we obtain

$$\begin{aligned} u \approx{}& \frac{540 + 473\epsilon\delta^2a^2}{198\epsilon} + \frac{86\delta^2ca^2}{3} \left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right] \left\{1 + c\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right]\right\} \\ & -\frac{1}{4}\left\{\frac{540 + 473\epsilon\delta^2a^2}{198\epsilon} + \frac{86\delta^2ca^2}{3} \left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right] \left\{1 + c\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right]\right\}\right\}^2 \\ & +\epsilon\delta^2\left(\frac{1}{3} - \frac{1}{2}z^2\right) u^*(\xi) \end{aligned} \tag{4.6}$$

where $u^*(\xi)$ is

$$u^* = \frac{86}{3}\delta^2ca^4\left[\frac{\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]} + 21c\frac{\exp[a(\xi + \xi_0)]^2}{\{1 - c\exp[a(\xi + \xi_0)]\}^2} + 12c^2\frac{\exp[a(\xi + \xi_0)]^3}{\{1 - c\exp[a(\xi + \xi_0)]\}^3} + 6c^3\frac{\exp[a(\xi + \xi_0)]^3}{\{1 - c\exp[a(\xi + \xi_0)]\}^4}\right] \tag{4.7}$$

Let now the equation of Riccati (3.9) be used as simplest equation. We again search solution of (3.16) in the form (3.2) with $\nu = 0$. For this case the balance equation is (3.19) The application of the modified method of simplest equation leads to a system of 8 algebraic relationships among the parameters of the extended Korteweg - de Vries equation and the parameters of the solution. One solution of this nonlinear system of algebraic relationships is

$$\begin{aligned}
\theta_2 &= -\frac{6a^2(p_5+2p_6)}{p_4}; \quad \theta_1 = -\frac{6ac(p_5+2p_6)}{p_4} \\
\theta_0 &= -\frac{3p_5p_6c^2+16p_6^2ad+8p_5^2ad+24p_5p_6ad+p_2p_5+2p_2p_6+p_5^2c^2+2p_6^2c^2-2p_3p_4}{2p_4(p_5+p_6)} \\
v &= \frac{v_1}{v_2} \\
v_1 &= \frac{1}{4}(-4p_2p_6p_3p_4-40p_5^2ac^2dp_6^2-16p_5ac^2dp_6^3+4p_5^3c^4p_6+5p_5^2c^4p_6^2+2p_5p_6^3c^4+ \\
&\quad 16p_5^4a^2d^2-32p_5^3ac^2dp_6-8p_5^4ac^2d+64p_5^3a^2d^2p_6+80p_5^2a^2d^2p_6^2+32p_5p_6^3a^2d^2+ \\
&\quad p_5^4c^4-2p_6p_2^2p_5-p_2^2p_5^2+4p_3^2p_4^2) \\
v_2 &= p_1p_4(p_5+p_6)^2
\end{aligned} \tag{4.8}$$

The solution of Eq. (3.16) based on the solution (3.10) of the equation of Riccati is

$$\eta = -\frac{3p_5p_6c^2+16p_6^2ad+8p_5^2ad+24p_5p_6ad+p_2p_5+2p_2p_6+p_5^2c^2+2p_6^2c^2-2p_3p_4}{2p_4(p_5+p_6)} - \\
\frac{6ac(p_5+2p_6)}{p_4}\left\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left[\frac{\theta(\xi+\xi_0)}{2}\right]\right\}\left\{1+\frac{a}{c}\left\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left[\frac{\theta(\xi+\xi_0)}{2}\right]\right\}\right\} \tag{4.9}$$

The solution of Eq. (3.16) based on the solution (3.11) of the equation of Riccati is

$$\begin{aligned}
\eta = &-\frac{3p_5p_6c^2+16p_6^2ad+8p_5^2ad+24p_5p_6ad+p_2p_5+2p_2p_6+p_5^2c^2+2p_6^2c^2-2p_3p_4}{2p_4(p_5+p_6)} - \\
&\frac{6ac(p_5+2p_6)}{p_4}\left\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)+\right. \\
&\left.\frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta}+2C^*\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\right\}\times \\
&\left\{1+\frac{a}{c}\left\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)+\right.\right. \\
&\left.\left.\frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta}+2C^*\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\right\}\right\}\right\}
\end{aligned} \tag{4.10}$$

We remember that in Eq. (4.9) and Eq. (4.10) $\theta^2 = c^2 - 4ad > 0$.

Taking into account the relationships (3.17) for the surface wave connected to the solution (4.9) is

$$\begin{aligned} u \approx & -\frac{16}{1089\epsilon^4\delta^4}\Bigg\{\Bigg(-\frac{15609}{64}\epsilon^5\delta^7\theta^4+\frac{223729}{256}\epsilon^4\delta^8\theta^4+\frac{46827}{128}\epsilon^5\delta^7\theta^4z^2\Bigg)\times \\ & \left[\tanh\left(\frac{1}{2}\theta(\xi+\xi_0)\right)\right]^4-\Bigg(\frac{21285}{32}\epsilon^3\delta^6\theta^2-\frac{15609}{32}\epsilon^5\delta^7\theta^4z^2+\frac{223729}{48}\epsilon^4\delta^8ad\theta^2- \\ & \frac{223729}{192}\epsilon^4\delta^8c^2\theta^2+\frac{5203}{16}\epsilon^5\delta^7\theta^4-\frac{15609}{32}\epsilon^4\delta^6\theta^2\Bigg)\left[\tanh\left(\frac{1}{2}\theta(\xi+\xi_0)\right)\right]^2- \\ & \Bigg(-\frac{1485}{8}\epsilon^3\delta^4+\frac{2025}{16}\epsilon^2\delta^4+\frac{223729}{576}\epsilon^4\delta^8c^4+\frac{5203}{16}\epsilon^4\delta^6c^2-\frac{223729}{72}\epsilon^4\delta^8adc^2- \\ & \frac{7095}{16}\epsilon^3\delta^6c^2+\frac{223729}{36}\epsilon^4\delta^8a^2d^2+\frac{15609}{128}\epsilon^5\delta^7\theta^4z^2-\frac{5203}{64}\epsilon^5\delta^7\theta^4+ \\ & \frac{7095}{4}\epsilon^3\delta^6ad-\frac{5203}{4}\epsilon^4\delta^6ad\Bigg)\Bigg\} \end{aligned} \tag{4.11}$$

Let now the extended tanh equation (3.12) be used as simplest equation. The balance equation is the same as in the case when the equation of Riccati is used as simplest equation. The application of the modified method of scimplest equation leads to a system of nonlinear algebraic relationships among the parameters of the equations and the parameters of the solution. One solution of this system is

$$\begin{aligned} \theta_2 &= -6\frac{\overline{a}^4(p_5+2p_6)}{p_4}; \qquad \theta_1 = 0 \\ \theta_0 &= \frac{-p_2p_5-2p_2p_6+24\overline{a}^2\overline{d}^2p_5p_6+16\overline{a}^2\overline{d}^2p_6^2+8\overline{a}^2\overline{d}^2p_5^2+2p_3p_4}{2p_4(p_5+p_6)} \\ v &= \frac{V}{4p_1p_4(p_5+p_6)^2} \\ V &= -2p_2^2p_5p_6-p_2^2p_5^2+16p_5^4d^4a^4+4p_3^2p_4^2+80p_5^2\overline{d}^4\overline{a}^4p_6^2+64p_5^3\overline{d}^4\overline{a}^4p_6+ \\ & \quad 32p_5\overline{d}^4\overline{a}^4p_6^3-4p_2p_3p_4p_6 \end{aligned} \tag{4.12}$$

and the solution of Eq. (3.16) is

$$\begin{aligned} \eta(\xi) = & \frac{-p_2p_5-2p_2p_6+24\overline{a}^2\overline{d}^2p_5p_6+16\overline{a}^2\overline{d}^2p_6^2+8\overline{a}^2\overline{d}^2p_5^2+2p_3p_4}{2p_4(p_5+p_6)}- \\ & 6\frac{\overline{a}^2\overline{d}^2(p_5+2p_6)}{p_4}\tanh^2[\overline{a}\overline{d}(\xi+\xi_0)] \end{aligned} \tag{4.13}$$

The surface wave (2.17) corresponding to the solution (4.13) is

$$\begin{aligned} u(\xi) \approx & \frac{1}{9801\epsilon^2}\big[\overline{a}^4\overline{d}^4\delta^4\epsilon^2(-561924\epsilon-2013561+842886\epsilon z^2)[\tanh(\overline{a}\overline{d}(\xi+\xi_0))]^4+ \\ & \overline{a}^2\overline{d}^2\delta^2\epsilon(-280962\epsilon-2684748a^2d^2\epsilon\delta^2+383130+749232\epsilon^2\delta^2a^2d^2- \\ & 1123848\epsilon^2\delta^2a^2d^2z^2)[\tanh(\overline{a}\overline{d}(\xi+\xi_0))]^2+26730\epsilon-187308a^2d^2\epsilon^2\delta^2- \\ & 894916a^4d^4\epsilon^2\delta^4-18225+255420a^2d^2\epsilon\delta^2-187308\epsilon^3\delta^4a^4d^4+ \\ & 280962\epsilon^3\delta^4a^4d^4z^2\big] \end{aligned} \tag{4.14}$$

For a further demonstration of the modified method of simplest equations let us now study what solutions of the classic Kortweg - de Vries equation can be obtained by this method. The classic Kortweg - de Vries equation is particular case of Eq. (3.16) for the case $p_4 = p_5 = p_6 = 0$. In other words we shall discuss the equation

$$p_1 \frac{\partial \eta}{\partial \tau} + p_2 \eta \frac{\partial \eta}{\partial \zeta} + p_3 \frac{\partial^3 \eta}{\partial \zeta^3} = 0 \tag{4.15}$$

which after the subsitution of $p_{1,2,3}$ from Eq. (3.17) is reduced to the Korteweg - de Vries equation. The balance equations for Eq. (4.15) are the same as for the Eq. (3.16). Lt us first discuss the case where the equation of Bernoulli is used as simplest equation. The application of the modified method of simplest equation reduces (4.15) to a system of 5 nonlinear algebraic relationships among the parameters of the equation and parameters of the solution (3.2) (we remember that $\nu = 0$ and $V(\xi)$) is a solution of the Bernoulli equation). A solution of this nonlinear algebraic system is

$$\theta_2 = -12 \frac{p_3 c^2}{p_2}; \quad \theta_1 = -12 \frac{p_3 c a}{p_2}; \quad \theta_0 = \frac{p_1 v - p_3 a^2}{p_2} \tag{4.16}$$

The corresponding solution of (4.15) for $c < 0$ and $a > 0$ is

$$\eta = \frac{p_1 v - p_3 a^2}{p_2} - 12 \frac{p_3 c a^2}{p_2} \left[\frac{\exp(a[\xi + \xi_0)]}{1 - c \exp[a(\xi + \xi_0)]} \right] \left\{ 1 + c \left[\frac{\exp(a[\xi + \xi_0)]}{1 - c \exp[a(\xi + \xi_0)]} \right] \right\} \tag{4.17}$$

For the case $a < 0$; $c > 0$ the solution of (4.15) is

$$\eta = \frac{p_1 v - p_3 a^2}{p_2} + 12 \frac{p_3 c a^2}{p_2} \left[\frac{\exp(a[\xi + \xi_0)]}{1 + c \exp[a(\xi + \xi_0)]} \right] \left\{ 1 - c \left[\frac{\exp(a[\xi + \xi_0)]}{1 + c \exp[a(\xi + \xi_0)]} \right] \right\} \tag{4.18}$$

By setting $p_1 = 2$; $p_2 = 3$ and $p_3 = \frac{1}{3}\delta^2$ in Eqs. (4.17) and (4.18) we obtain the corresponding solutions of the classic Korteweg - de Vries equation.

Let now the equation of Riccati (3.9) is used as simplest equation. For this case the application of the modified method of simplest equation leads to a system of 6 nonlinear relationships among the parameters of the equation and parameters of the solution. One solution of this system is

$$\theta_2 = -12 \frac{p_3 a^2}{p_2}; \quad \theta_1 = -12 \frac{p_3 a c}{p_2}; \quad \theta_0 = \frac{p_1 v - 8 p_3 a d - p_3 c^2}{p_2} \tag{4.19}$$

The solution of Eq. (4.15) based on the solution (3.10) of the Riccati equation is

$$\begin{aligned} \eta \;\; = \;\; & \frac{p_1 v - 8 p_3 a d - p_3 c^2}{p_2} + \frac{p_3 a c}{p_2} \left\{ \frac{c}{2a} + \frac{\theta}{2a} \tanh \left[\frac{\theta(\xi + \xi_0)}{2} \right] \right\} \left\{ 1 - \right. \\ & \left. \frac{a}{c} \left\{ \frac{c}{2a} + \frac{\theta}{2a} \tanh \left[\frac{\theta(\xi + \xi_0)}{2} \right] \right\} \right\} \end{aligned} \tag{4.20}$$

and the solution of Eq. (4.15) based on the solution (3.11) of the Riccati equation is

$$\eta = \frac{p_1 v - 8p_3 a d - p_3 c^2}{p_2} - 12\frac{p_3 a c}{p_2}\Bigg\{ -\frac{c}{2a} - \frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right) + \frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta} + 2C^* \exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\Bigg\} + \Bigg\{1 + \frac{a}{c}\Bigg[-\frac{c}{2a} - \frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right) + \frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta} + 2C^* \exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\Bigg]\Bigg\} \tag{4.21}$$

From Eqs. (4.20) and (4.21) by setting $p_1 = 2$; $p_2 = 3$ and $p_3 = \frac{1}{3}\delta^2$ one obtains the corresponding solutions of the classic Korteweg - de Vries equation.

Finally the the extended tahn equation (3.12) be used as a simplest equation. The balance equation for this case is the same as in the case when the simplest equation is the equation of Riccati. The application of the modified method of simplest equation leads to a system of 6 nonlinear algebraic equations for the parameters of the solution. One solution of this system is

$$\theta_2 = -12\frac{p_3\overline{a}^4}{p_2}; \quad \theta_1 = 0; \quad \theta_0 = \frac{vp_1 + 8p_3\overline{a}^2\overline{d}^2}{p_2} \tag{4.22}$$

The sollution of (4.15) is

$$\eta(\xi) = \frac{vp_1 + 8p_3\overline{a}^2\overline{d}^2}{p_2} - 12\frac{p_3\overline{a}^4}{p_2}\frac{\overline{d}^2}{\overline{a}^2}\tanh^2[\overline{a}\overline{d}(\xi+\xi_0)] \tag{4.23}$$

5 Exact traveling - wave solutions of the generalized Camassa - Holm equation

Let first the equation of Bernoulli be used as simplest equation and the values of parameters from (3.20) are $J = 3$, $I = 2$, $k = 2$. Then from (3.20) one obtains $\nu_1 = 2$. The equation we shall solve is obtained from (2.20) when $I = 2$, $J = 3$. This equation is

$$\frac{\partial U}{\partial T} + p_1\frac{\partial U}{\partial X} - p_2\frac{\partial^3 U}{\partial X^2 \partial T} - p_3\frac{\partial^3}{\partial X^3}(q_0 + q_1 U + q_2 U^2) + p_4\frac{\partial}{\partial X}\left[\frac{1}{2}(r_0 + r_1 U + r_2 U^2 + r_3 U^3) + p_5 q_2\left(\frac{\partial U}{\partial X}\right)^2\right] = 0 \tag{5.1}$$

The solution of Eq. (2.21) is searched in the form

$$U(\xi) = \theta_0 + \theta_1 V(\xi) + \theta_2 V(\xi)^2 \tag{5.2}$$

where $V(\xi)$ is solution of the equation of Bernoulli (3.6) for $k = 2$. The substitution of (5.2) in (2.21) leads to a large system of 7 nonlinear relationships among the parameters of the generalized Camassa - Holm equation and parameters of the solution (5.2) One solution of this system of equations is

$$\begin{aligned}
\theta_2 &= 8\frac{q_2 ac(5p_3 - p_4 p_5)}{p_4 r_3}; \quad \theta_1 = 8\frac{q_2 c^2(5p_3 - p_4 p_5)}{p_4 r_3} \\
\theta_0 &= \frac{\Theta_0}{6q_2 p_4 r_3(4p_3 - p_4 p_5)} \quad r_1 = \frac{R_1}{12p_2 p_4 r_3(4p_3 - p_4 p_5)} \\
\Theta_0 &= -36q_2^2 a^2 p_4 p_5 p_3 + 4q_2^2 a^2 p_4^2 p_5^2 - 3p_2 v p_4 r_3 + 3p_3 q_1 p_4 r_3 + 80q_2^2 a^2 p_3^2 - \\
&\quad 10q_2 p_3 r_2 p_4 + 2q_2 p_4^2 p_5 r_2 \\
R_1 &= -(1520q_2^4 a^4 p_4^2 p_5^2 p_3^2 - 256q_2^4 a^4 p_4^3 p_5^3 p_3 - 4q_2^2 p_4^4 p_5^2 r_2^2 + 32q_2^2 p_3 r_2^2 p_4^3 p_5 - \\
&\quad 60q_2^2 p_3^2 r_2^2 p_4^2 + 3840q_2^4 a^4 p_3^4 - 12p_3^2 q_1 p_4^2 r_3 q_2 r_2 + 12p_2 v p_4^2 r_3 q_2 p_3 r_2 - \\
&\quad 18p_2 v p_4^2 r_3^2 p_3 q_1 + 9p_3^2 q_1^2 p_4^2 r_3^2 + 16q_2^4 a^4 p_4^4 p_5^4 + 9p_2^2 v^2 p_4^2 r_3^2 - \\
&\quad 3968q_2^4 a^4 p_4 p_5 p_3^3 - 24v q_2^2 r_3 p_4^3 p_5^2 + 192v q_2^2 r_3 p_4^2 p_3 p_5 - 384v q_2^2 r_3 p_4 p_3^2 - \\
&\quad 192p_1 q_2^2 r_3 p_4^2 p_3 p_5 + 384p_1 q_2^2 r_3 p_4 p_3^2 + 24p_1 q_2^2 r_3 p_4^3 p_5^2)
\end{aligned} \tag{5.3}$$

The corresponding solution of Eq. (5.1) is

$$\begin{aligned}
U(\xi) = \frac{\Theta_0}{6q_2 p_4 r_3(4p_3 - p_4 p_5)} &+ 8\frac{q_2 c^2(5p_3 - p_4 p_5)}{p_4 r_3}\left\{\frac{a\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right\} + \\
&8\frac{q_2 ac(5p_3 - p_4 p_5)}{p_4 r_3}\left\{\frac{a\exp[a(\xi + \xi_0)]}{1 - c\exp[a(\xi + \xi_0)]}\right\}^2
\end{aligned} \tag{5.4}$$

for $c < 0$ and $a > 0$. For $c > 0$ and $a < 0$ the solution of Eq. (5.1) is

$$\begin{aligned}
U(\xi) = \frac{\Theta_0}{6q_2 p_4 r_3(4p_3 - p_4 p_5)} &- 8\frac{q_2 c^2(5p_3 - p_4 p_5)}{p_4 r_3}\left\{\frac{a\exp[a(\xi + \xi_0)]}{1 + c\exp[a(\xi + \xi_0)]}\right\} + \\
&8\frac{q_2 ac(5p_3 - p_4 p_5)}{p_4 r_3}\left\{\frac{a\exp[a(\xi + \xi_0)]}{1 + c\exp[a(\xi + \xi_0)]}\right\}^2
\end{aligned} \tag{5.5}$$

Now let the equation of Riccati (3.9) be used as simplest equation. From the possible balance equations Eq. (3.23) will be used below with values of the parameters $J = 1$; $\nu_1 = 2$; $I = 2$. Thus the equation which will be solved is

$$\begin{aligned}
\frac{\partial U}{\partial T} + p_1\frac{\partial U}{\partial X} - p_2\frac{\partial^3 U}{\partial X^2 \partial T} - p_3\frac{\partial^3}{\partial X^3}(q_0 + q_1 U + q_2 U^2) + \\
p_4\frac{\partial}{\partial X}\left[\frac{1}{2}(r_0 + r_1 U) + p_5 q_2\left(\frac{\partial U}{\partial X}\right)^2\right] = 0
\end{aligned} \tag{5.6}$$

In this case the application of the method of simplest equation to Eq. (5.6) leads to a system of 8 nonlinear algebraic relationships among the parameters of the solved equation and the parameters of the solution. One solution of this system is

$$\begin{aligned}
\theta_2 &= -15\frac{a^2(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)} \\
\theta_1 &= -15\frac{ac(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)} \\
\theta_0 &= \frac{\Theta_0}{q_2p_4p_5(-8c^2da+16a^2d^2+c^4)} \\
\Theta_0 &= \frac{1}{4}(40adp_4r_1-80adv+80adp_1+5c^2p_4r_1-10c^2v+10c^2p_1+ \\
&\quad 16p_4p_5q_1c^2da-32p_4p_5q_1a^2d^2-2p_4p_5q_1c^4-80p_2vc^2da+160p_2va^2d^2+ \\
&\quad 10p_2vc^4)
\end{aligned} \tag{5.7}$$

The corresponding solution of Eq. (5.6) based on the solution (3.10) of the Riccati equation is

$$\begin{aligned}
U(\xi) &= \frac{\Theta_0}{q_2p_4p_5(-8c^2da+16a^2d^2+c^4)}+ \\
&\quad 15\frac{ac(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)}\left[\frac{c}{2a}+\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]- \\
&\quad 15\frac{a^2(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)}\left[\frac{c}{2a}+\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]^2
\end{aligned} \tag{5.8}$$

where $\theta^2 = c^2-4ad > 0$.

The solution of (5.6) based on the solution (3.11) of the Riccati equation is

$$\begin{aligned}
U(\xi) &= \frac{\Theta_0}{q_2p_4p_5(-8c^2da+16a^2d^2+c^4)}- \\
&\quad 15\frac{ac(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)}\Bigg\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)+ \\
&\quad \frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta}+2C^*\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\Bigg\}- \\
&\quad 15\frac{a^2(-p_4r_1+2v-2p_1)}{q_2p_4p_4(-8c^2da+16a^2d^2+c^4)}\Bigg\{-\frac{c}{2a}-\frac{\theta}{2a}\tanh\left(\frac{\theta(\xi+\xi_0)}{2}\right)+ \\
&\quad \frac{\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)}{2\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\left[\frac{a}{\theta}+2C^*\exp\left(\frac{\theta(\xi+\xi_0)}{2}\right)\cosh\left(\frac{\theta(\xi+\xi_0)}{2}\right)\right]}\Bigg\}^2
\end{aligned} \tag{5.9}$$

Let now the extended tanh equation (3.12) be the simplest equation. Let in addition $I = J = 2$ and $\nu_1 = 2$. The equation which is solved for this case is

$$\begin{aligned} &\frac{\partial U}{\partial T} + p_1 \frac{\partial U}{\partial X} - p_2 \frac{\partial^3 U}{\partial X^2 \partial T} - p_3 \frac{\partial^3}{\partial X^3}(q_0 + q_1 U + q_2 U^2) + \\ &\qquad p_4 \frac{\partial}{\partial X}\left[\frac{1}{2}(r_0 + r_1 U + r_2 U^2) + p_5 q_2 \left(\frac{\partial U}{\partial X}\right)^2\right] = 0 \end{aligned} \tag{5.10}$$

The application of the modified method of simplest equation leads to a system of 8 noninear algebraic relationships among the parameters of the solution and para meters of the solved equation. One solution of this system is

$$\begin{aligned} \theta_2 &= 60\overline{a}^4 \frac{(-2vp_5q_2 + 2p_1p_5q_2 + r_1p_4p_5q_2 + 5r_2p_2v - r_2p_4p_5q_1)}{p_4(64p_5^2q_2^2\overline{d}^4\overline{a}^2 - 25r_2^2)} \\ \theta_1 &= 0 \\ \theta_0 &= \frac{\Theta_0}{p_4(-25r_2^2 + 64p_5^2q_2^2\overline{d}^4\overline{a}^4)} \\ \Theta_0 &= \frac{1}{2}(160p_5q_2\overline{a}^2\overline{d}^2v - 160p_5q_2\overline{a}^2\overline{d}^2p_1 - 80p_5q_2\overline{a}^2\overline{d}^2r_1p_4 - 400\overline{a}^2\overline{d}^2r_2p_2v + \\ &\quad 80p_5\overline{a}^2\overline{d}^2r_2p_4q_1 + 320p_2vp_5q_2\overline{d}^4\overline{a}^4 - 50r_2v + 50r_2p_1 + 25r_2r_1p_4 - \\ &\quad 64p_4p_5^2q_1q_2\overline{d}^4\overline{a}^4) \\ p_3 &= \frac{1}{5}p_4p_5 \end{aligned} \tag{5.11}$$

The corresponding solution is as follows

$$\begin{aligned} U(\xi) &= \frac{\Theta_0}{p_4(-25r_2^2 + 64p_5^2q_2^2\overline{d}^4\overline{a}^4)} + \\ &\quad 60\overline{a}^2\overline{d}^2 \frac{(-2vp_5q_2 + 2p_1p_5q_2 + r_1p_4p_5q_2 + 5r_2p_2v - r_2p_4p_5q_1)}{p_4(64p_5^2q_2^2\overline{d}^4\overline{a}^2 - 25r_2^2)} \tanh^2[\overline{a}\overline{d}(\xi + \xi_0)] \end{aligned} \tag{5.12}$$

Let us now discuss the equation

$$\frac{\partial U}{\partial T} + p_1 \frac{\partial U}{\partial X} - p_2 \frac{\partial^3 U}{\partial X^2 \partial T} + 3p_4 U \frac{\partial U}{\partial X} - 2p_3 \frac{\partial U}{\partial X}\frac{\partial^2 U}{\partial X^2} - p_5 U \frac{\partial^3 U}{\partial X^3} = 0 \tag{5.13}$$

which can be reduced to Eq. (2.21) and then to the Camassa - Holm equation when $p_5 = p_3$. If the equation of Bernoulli (3.6) is used as simplest equation in the modified method of simplest equation we obtain balance equation of the kind $2\nu_1 + 3(k-1) = 2\nu_1 + 3(k-1)$. The simplest case is $\nu_1 = 2$, $k = 2$. For this case the application of the modified method of simplest equation reduces Eq. (5.13) to a system of 7 nonlinear algebraic relationships among the parsmeters of the equation and the parameters of the solution. One solution of this system is

$$\begin{aligned} p_5 &= -p_3; \qquad v = \frac{12p_1c^2p_3 - 9p_4^2\theta_2 + p_3^2\theta_2a^4}{12c^2(p_3+3p_4p_2)} \\ \theta_0 &= -\frac{3p_4\theta_2 - p_3\theta_2a^2 - 3\theta_2a^2p_4p_2 + 12p_2p_1c^2 + p_2p_3\theta_2a^4}{12c^2(p_3+3p_4p_2)} \\ \theta_1 &= \frac{\theta_2 a}{c} \end{aligned} \tag{5.14}$$

the corresponding solution of Eq. (5.13) is

$$\begin{aligned} U(\xi) = -&\frac{3p_4\theta_2 - p_3\theta_2a^2 - 3\theta_2a^2p_4p_2 + 12p_2p_1c^2 + p_2p_3\theta_2a^4}{12c^2(p_3+3p_4p_2)} + \\ &\frac{\theta_2 a}{c}\left\{\frac{a\exp[a(\xi+\xi_0)]}{1-c\exp[a(\xi+\xi_0)]}\right\} + \theta_2\left\{\frac{a\exp[a(\xi+\xi_0)]}{1-c\exp[a(\xi+\xi_0)]}\right\}^2 \end{aligned} \tag{5.15}$$

for the case $c < 0$, $a > 0$. For the case $c > 0$, $a < 0$ the solution is

$$\begin{aligned} U(\xi) = -&\frac{3p_4\theta_2 - p_3\theta_2a^2 - 3\theta_2a^2p_4p_2 + 12p_2p_1c^2 + p_2p_3\theta_2a^4}{12c^2(p_3+3p_4p_2)} - \\ &\frac{\theta_2 a}{c}\left\{\frac{a\exp[a(\xi+\xi_0)]}{1+c\exp[a(\xi+\xi_0)]}\right\} + \theta_2\left\{\frac{a\exp[a(\xi+\xi_0)]}{1+c\exp[a(\xi+\xi_0)]}\right\}^2 \end{aligned} \tag{5.16}$$

It must be noted that because of the requirement $p_5 = -p_3$ the above two solutions can not be reduced to solutions of the Camassa - Holm equation. If the rquirement was $p_5 = p_3$ such reduction would be possible. However the requirement is $p_5 = -p_3$ and the reduction is not possible.

6 Concluding remarks

In this paper we have studied two nonlinear partial differential equations connected to the water waves: extended Korteweg - de Vries equation and generalized Camassa - Holm equation. The goal of the study was to obtain exact traveling-wave solutions of these equations. This goals was achieved on the basis of the method of simplest equation which is an effective tool for obtaining exact analytical solutions for large classes of nonlinear partial differential equations. We have applied a modification of this method based on use of balance equation. The equations of Bernoulli, Riccati, and extended tanh - equation have been used as simplest equations. We have obtained numerous exact traveling-wave solutions of the studied equations and for the case of the extended Korteweg - de Vries equation we have obtained three expressions for surface water waves which correspond to the exact solutions of the extended Korteweg - de Vries equation. As a side result we have applied the methodology to the classic Korteweg - de Vries equation and we have demonstrated that by means of the modified method of simplest equation exact solutions of this classic nonlinear partial differential equation can be obtained too.

This paper is manuscript Nr. AMC-D-10-04363 submitted to *Applied Mathematics and Computation* on 27th of December 2010. The manuscript is still under review.

References

[1] J. C. Tannenhill, D. A. Anderson, R. H. Pletcher. Computational fluid mechanics and heat transfer. Taylor and Francis, Philadelphia, PA, 1997.

[2] F. Dias. Nonlinear gravity and capillary-gravity waves. Annual Review of Fluid Mechanics 31 (1999) 301 - 346.

[3] N. K. Vitanov, F. H. Busse. Bounds on the heat transport in a horizontal fluid layer with stress-free boundaries. ZAMP 48 (1997) 310 - 324.

[4] A. C. Scott. Nonlinear science. Emergence and dynamics of coherent structures. Oxford University Press, Oxford, UK, 1999.

[5] N. K. Vitanov. Upper bound on the heat transport in a horizontal fluid layer of infinite Prandtl number. Phys. Lett. A 248 (1998) 338 - 346.

[6] T. B. Benjamin, J. L. Bona, J. J. Mahoni. Model equations for long waves in nonlinear dispersive systems. Phil. Trans. Roy. Soc. London A 272 (1972) 47 - 78.

[7] J. W. Miles. Solitary waves. Annual Review of Fluid Mechanics 12 (1980) 11 - 43.

[8] H. Kantz, D. Holstein, M. Ragwitz, N. K. Vitanov. Markov chain model for turbulent wind speed data. Physica A 342 (2004) 315 - 321.

[9] N. K. Vitanov. Upper bound on the heat transport in a layer of fluid of infinite Prandtl number, rigid lower boundary, and stress-free upper boundary. Phys. Rev. E 61 (2000) 956 - 959.

[10] A. Scott. Neuroscience: A mathematical primer. Springer, New York, 2002.

[11] A. Longtin, J. G. Milton. Modeling autonomous oscillations in the human pupil light reflex using nonlinear delay-differential equations. Bull. Math. Biology 51 (1989) 605 - 624.

[12] M. Ablowitz, P. A. Clarkson. Solitons, nonlinear evolution equations and inverse scattering. Cambridge University Press, Cambridge, UK, 1991.

[13] A. -M. Wazwaz. Partial differential equations and solitary waves theory. Springer, Berlin, 2009.

[14] E. Infeld, G. Rowlands. Nonlinear waves, solitons and chaos. Cambridge University Press, Cambridge, UK, 1990.

[15] A. C. Newell. Solitons in mathematics and physics. SIAM, Philadelphia, PA, 1985.

[16] J. D. Murray. Lectures on nonlinear differential equation models in biology. Oxford University Press, Oxford, England, 1977.

[17] L. Perko. Differential equations and dynamical systems. Springer, New York, 2001.

[18] R. M. May. Stability and complexity in model ecosystems. Princeton University Press, New Jersey, 2001.

[19] R. Temam. Navier - Stokes equations: Theory and numerical analysis. AMS Chelsea Publishing, Providence, R. I., 2001.

[20] C. Foias, O. Manley, R. Rosa, R. Temam. Navier - Stokes equations and turbulence. Cambridge University Press, Cambridge, UK, 2001.

[21] P. Holmes, J. L. Lumley, G. Berkooz. Turbulence, coherent structures, dynamical systems and symmetry. Cambridge University Press, Cambridge, UK, 1996.

[22] N. P. Hoffmann, N. K. Vitanov. Upper bounds on energy dissipation in Couette-Ekman flow. Phys. Lett. A 255 (1999) 277 - 286.

[23] N. K. Vitanov. Upper bounds on the heat transport in a porous layer. Physica D 136 (2000) 322 - 339.

[24] T. Boeck, N. K. Vitanov. Low-dimensional chaos in zero-Prandtl-number Benard-Marangoni convection. Phys. Rev. E 65 (2002) Article number 037203.

[25] N. K. Vitanov. Convective heat transport in a fluid layer of infinite Prandtl number: upper bounds for the case of rigid lower boundary and stress - free upper boundary. European Physical Journal B 15 (2000) 349 - 355.

[26] N. K. Vitanov. Upper bounds on convective heat transport in a rotating layer of infinite Prandtl number: Case of intermediate taylor numbers. Phys. Rev. E 62 (2000) 3581 - 3591.

[27] C. S. Gardner, J. M. Greene, M. D. Kruskal, R. R. Miura. Method for solving Korteweg- de Vries equation. Phys. Rev. Lett. 19 (1967) 1095 - 1097.

[28] M. J. Ablowitz, D. J. Kaup, A. C. Newell. Nonlinear evolution equations of physical significance. Phys. Rev. Lett. 31 (1973) 125 - 127.

[29] M. Remoissenet. Waves called solitons. Springer, Berlin; 1993.

[30] M. J. Ablowitz, D. J. Kaup, A. C. Newell, H. Segur. Inverse scattering transform - Fourier analysis for nonlinear problems. Studies in Applied Mathematics 53 (1974) 249 - 315.

[31] R. Hirota. Exact solution of Korteweg-de Vries equation for multiple collisions of solitons. Phys. Rev. Lett. 27 (1971) 1192 - 1194.

[32] N. A. Kudryashov. Exact solutions of the generalized Kuramoto - Sivashinsky equation. Phys. Lett. A 147 (1990) 287 - 291.

[33] Z. Y. Yan. New explicit travelling wave solutions for two new integrable coupled nonlinear evolution equations. Phys. Lett. A 292 (2001) 100 - 106.

[34] E. G. Fan. Extended tanh-function method and its application to nonlinear equations. Phys. Lett. A 277 (2000) 212 - 218.

[35] E. G. Fan. Traveling wave solutions for nonlinear equations using symbolic computation. Computers & Mathematics with Applications 43 (2002) 671 - 680.

[36] J. -H. He, X,-H. Wu. Exp-function method for nonlinear wave equations. Chaos Solitons & Fractals 30 (2006) 700 - 708.

[37] N. A. Kudryashov. Simplest equation method to look for exact solutions of nonlinear differential equations. Chaos Solitons & Fractals 24 (2005) 1217 - 1231.

[38] N. K. Vitanov, Z. I. Dimitrova, H. Kantz. Modified method of simplest equation and its application to nonlinear PDEs. Applied Mathematics and Computation 216 (2010) 2587 - 2595.

[39] N. K. Vitanov. Modified method of simplest equation: Powerful tool for obtaining exact and approximate travelling-wave solutions of nonlinear PDEs. Communications in Nonlinear Science and Numerical Simulation 16 (2011) 1176 - 1185.

[40] M. J. Ablowitz, A. Zeppetela. Explicit solutions of Fisher equation for a specifical wave speed. Bull. Math. Biol. 41 (1979) 835 - 840.

[41] Z. I. Dimitrova, N. K. Vitanov. Influence of adaptation on the nonlinear dynamics of a system of competing populations. Phys. Lett. A 272 (2000) 368 - 380.

[42] Z. I. Dimitrova, N. K. Vitanov. Dynamical consequences of adaptation of growth rates in a system of three competing populations. J. Phys. A: Math. Gen. 34 (2001) 7459 - 7473.

[43] Z. I. Dimitrova, N. K. Vitanov. Adaptation and its impact on the dynamics of a system of three competing populations. Physica A 300 (2001) 91 - 115.

[44] Z. I. Dimitrova, N. K. Vitanov. Chaotic pairwise competition. Theoretical Population Biology 66 (2004) 1 - 12.

[45] N. K. Vitanov, Z. I. Dimitrova, H. Kantz. On the trap of extinction and its elimination. Phys. Lett. A 349 (2006) 350 - 355.

[46] N. K. Vitanov, I. P. Jordanov, Z. I. Dimitrova. On nonlinear dynamics of interacting populations: Coupled kink waves in a system of two populations. Communications in Nonlinear Science and Numerical Simulation 14 (2009) 2379 - 2388.

[47] N. K. Vitanov, I. P. Jordanov, Z. I. Dimitrova. On nonlinear population waves. Applied Mathematics and Computation. 215 (2009) 2950 - 2964.

[48] X. Y. Wang. Exact and explicit wave solutions for the generalized Fisher equation. Phys. Lett. A 131 (1988) 277 - 279.

[49] N. A. Kudryashov. Exact solitary waves of the Fisher equation. Phys. Lett. A 342 (2005) 99 - 106.

[50] N. K. Vitanov, Z. I. Dimitrova. Application of the method of simplest equation for obtaining exact traveling-wave solutions for two classes of model PDEs from ecology and population dynamics. Communications in Nonlinear Science and Numerical Simulation 15 (2010) 2836 - 2845.

[51] E. E. Holmes, M. A. Lewis, J. E. Banks, R. R. Veit. Partial differential equation in ecology: Spatial interactions and population dynamics. Ecology 75 (1994) 17 - 29.

[52] N. K. Vitanov, Z. I. Dimitrova, M. Ausloos. Verhulst - Lotka - Volterra (VLV) model of ideological struggle. Physica A 389 (2010) 4970 - 4980.

[53] S. Lou. Symmetry analysis and exact solutions of the 2+1 - dimensional sine - Gordon system. J. Math. Phys. 41 (2000) 6509 - 6524.

[54] A. -M. Wazwaz. The tanh method: exact solutions of the sine - Gordon and the sh-Gordon equations. Applied Mathematics and Computation 167 (2005) 1196 - 1210.

[55] N. K. Martinov, N. K. Vitanov. New class of running-wave solutions of the 2+1-dimensional sine-Gordon equation. J. Phys. A: Math. Gen. 27 (1994) 4611 - 4618.

[56] N. K. Vitanov. On traveling waves and double - periodic structures in two - dimensional sine - Gordon systems. J. Phys. A: Math. Gen. 29 (1996) 5195 - 5207.

[57] P. A. Clarkson, E. L. Mansfield, A. E. Milne. Symmetries and exact solutions of a (2+1)-dimensional sine - Gordon system. Phil. Trans. Roy. Soc. London A 354 (1996) 1807 - 1835.

[58] N. K. Vitanov, N. K. Martinov. On the solitary waves in the sine-Gordon model of the two-dimensional Josephson junction. Z. Phys. B 100 (1996) 129 - 135.

[59] N. K. Vitanov. Breather and soliton wave families for the sine - Gordon equation. Proc. Roy. Soc. London A 454 (1998) 2409 - 2423.

[60] R. Radha, M. Lakshamanan. The (2+1) - dimensional sine - Gordon equation; Integrability and localized solutions. J. Phys A: Math. Gen. 29 (1996) 1551 - 1562.

[61] A. Nakamura. Exact cylindrical soliton solutions of the sine - Gordon equation, the sinh - Gordon equation and the periodic Toda equation. J. Phys. Society Japan 57 (1988) 3309 - 3322.

[62] N. K. Martinov, N. K. Vitanov. On the self - consistent thermal equilibrium structures in two - dimensional negative temperature systems. Canadian Journal of Physics 72 (1994) 618 - 624.

[63] A. M. Wazwaz. Exact solutions for the generalized sine - Gordon and sinh-Gordon equation. Chaos Solitons & Fractals 28 (2006) 127 - 135.

[64] S. Panchev, T. Spassova, N. K. Vitanov. Analytical and numerical investigation of two families of Lorenz-like dynamical systems. Chaos Solitons & Fractals 33 (2007) 1658 - 1671.

[65] N. A. Kudryashov, N. B. Loguinova. Extended simplest equation method for nonlinear differential equations. Applied Mathematics and Computation 205 (2008) 396 - 402.

[66] N. A. Kudryashov, M. V. Demina. Polygons of differential equations for finding exact solutions. Chaos Solitons & Fractals 33 (2007) 480 - 496.

[67] N. K. Vitanov. Application of simplest equations of Bernoulli and Riccati kind for obtaining exact traveling-wave solutions for a class of PDEs with polynomial nonlinearity. Communications in Nonlinear Science and Numerical Simulation 15 (2010) 2050 - 2060.

[68] N. K. Vitanov, Z. I. Dimitrova, K. N. Vitanov. On the class of nonlinear PDEs that can be treated by the modified method of simplest equation. Application to generalized Degasperis - Processi equation and b-equation. Communications in Nonliner Science and Numerical Simulation (in press), doi:10.1016/j.cnsns.2010.11.013

[69] D. J. Benney. Long nonlinear waves in fluid flows. Journal of Mathematics and Physics 45 (1966) 52 - 63.

[70] J. J. Stoker. Water waves. the mathematical theory with applications. Wiley, New York, 1992.

[71] R. S. Johnson. A modern introduction to the mathematical theory of water waves. Cambridge University Press, Cambridge, UK, 1997.

[72] R. S. Johnson. The classical problem of water waves: a reservoir of integrable and nearly integrable equations. Journal of Nonlinear Mathematical Physics 10 (2003) Supplement 1, 72 - 92.

[73] R. Camassa, D. D. Holm. An integrable shallow water equation with peaked solitons. Phys. Rev. Lett. 71 (1993) 1661 - 1664.

[74] L. Tian, Y. Wang. Global conservative solutions of the generalized Camassa - Holm equation. International Journal of Nonlinear Science 5 (2008) 195 - 202.

[75] L. Tian, X. Song. New peaked solitary wave solutions of the generalized Camassa-Holm equation. Chaos Solitons & Fractals 19 (2004) 621 - 637.

[76] L. Tian, J. Yin. New compacton solutions and solitary wave solutions of fully nonlinear generalized Camassa-Holm equations. Chaos Solitons & Fractals 20 (2004) 289 - 299.

[77] N. A. Kudryashov, N. B. Loguinova. Be careful with the exp - function method. Commun. Nonlinear Sci. Numer. Simulat. 14 (2009) 1881 - 1890.

[78] N. A. Kudryashov. Seven common errors in finding exact solutions of nonlinear differential equations. Commun. Nonlinear Sci. Numer. Simulat. 14 (2009) 3507 - 3529.